

Reconstructing 3-colored grids from horizontal and vertical projections is NP-hard

Christoph Dürr*

Flavio Guíñez†

Martín Matamala‡

April 21, 2009

Abstract

We consider the problem of coloring a grid using k colors with the restriction that in each row and each column has a specific number of cells of each color. In an already classical result, Ryser obtained a necessary and sufficient condition for the existence of such a coloring when two colors are considered. This characterization yields a linear time algorithm for constructing such a coloring when it exists. Gardner *et al.* showed that for $k \geq 7$ the problem is NP-hard. Afterward Chrobak and Dürr improved this result, by proving that it remains NP-hard for $k \geq 4$. We solve the gap by showing that for 3 colors the problem is already NP-hard. Besides we also give some results on tiling tomography problems.

1 Introduction

Tomography consists of reconstructing spatial objects from lower dimensional projections, and has medical applications as well as non-destructive quality control. In the discrete variant, the objects to be reconstructed are discrete, as for example atoms in a crystalline structure, see [1].

One of the first studied problem in discrete tomography involves the coloring of a grid, with a fixed number of colors with the requirement that each row and each column has a specific total number of entries of each color.

More formally we are given a set of colors $\mathcal{C}$, and an $m \times n$ matrix M , whose items are elements of $\mathcal{C}$. The projection of M is a sequence of vectors $r^c \in \mathbb{N}^m, s^c \in \mathbb{N}^n$, for $c \in \mathcal{C}$, where

$$r_i^c = |\{j : M_{ij} = c\}|, \quad s_j^c = |\{i : M_{ij} = c\}|.$$

In the reconstruction problem, we are given only a sequence of vectors satisfying for $1 \leq i \leq m$, $1 \leq j \leq n$, $c \in \mathcal{C}$,

$$\sum_c r_i^c = n, \quad \sum_c s_j^c = m, \quad \sum_i r_i^c = \sum_j s_j^c, \quad (1)$$

and the goal is to compute a matrix M that has the given projections. If there are $k = |\mathcal{C}|$ colors, we call it the k -COLOR TOMOGRAPHY PROBLEM.

It was known since long time, that for 2 colors, the problem can be solved in polynomial time [8]. Ten years ago it was shown that the problem is NP-hard for 7 colors [5]. By NP-hardness, we mean that the decision variant — deciding whether a given instance is *feasible*, i.e. admits a solution — is NP-hard. Shortly after this proof was improved to show NP-hardness for 4 colors, leaving open the case when $|\mathcal{C}| = 3$ [2]. This paper closes the gap, by showing that for 3 colors already the problem is NP-hard.

*CNRS, LIX (UMR 7161), Ecole Polytechnique, 91128 Palaiseau, France.

†DIM, Universidad de Chile, Casilla 170-3, Correo 3, Santiago, Chile.

‡DIM and CMM (UMI 2807, CNRS), Universidad de Chile, Casilla 170-3, Correo 3, Santiago, Chile.

Just to fix the notation, for $|\mathcal{C}| = 2$ we denote the colors as *black* and *white*, and use symbols B, W . For $|\mathcal{C}| = 3$ we denote the colors as *red*, *green* and *yellow* and use symbols R, G, Y . Notice that we can think white and yellow as ground colors in the 2 and 3-color problem, respectively. Thus when we denote the instance of the tomography problem, we sometimes omit the white or yellow projections as they are redundant. In addition for a 2-color instance (r^B, s^B) we omit the superscript when the context permits it.

First we recall some well known facts about the 2-COLOR TOMOGRAPHY PROBLEM.

Lemma 1 ([8]) *Let (r, s) be a feasible instance of the 2-color tomography problem. Let I be some set of rows, and J be some set of columns. If*

$$\sum_{i \in I} r_i - \sum_{j \notin J} s_j = |I \times J|, \quad (2)$$

then every solution to the instance will be all black in $I \times J$ and all white in $\bar{I} \times \bar{J}$.

Proof: The sets I, J divide the grid into four parts, $I \times J$, $I \times \bar{J}$, $\bar{I} \times J$ and $\bar{I} \times \bar{J}$. The value $\sum_{i \in I} r_i$ equals the number of black cells in the first two parts, and $\sum_{j \notin J} s_j$ the number of black cells in the second and last part. So the difference is the number of black cells in $I \times J$ minus the number of black cells in $\bar{I} \times \bar{J}$. So when (2) holds, the first part must be all black and the last part all white. $\square$

Before stating the next lemma, we need to introduce some notation about vectors. The *conjugate* of a vector $s \in \{0, 1, \dots, m\}^n$ is defined as the vector $s^* \in \{0, 1, \dots, n\}^m$ where $s_i^* = |\{j : s_j \geq i\}|$. There is a very simple graphical interpretation of this. Let be an $m \times n$ matrix M , such in column j , the first s_j cells are colored black and the others are colored white. Then the conjugate of s is just the row projection of M , see figure 1.




Figure 1: Example of a vector s and its conjugate s^* .

Note that s^* is always a non-increasing vector. If in addition s is non-increasing we have that $(s^*)^* = s$ since in this case $s_i^* = \max\{j : s_j \geq i\}$ and $s_i^* \geq j$ if and only if $s_j \geq i$.

For every $s, t \in \mathbb{N}^n$ we say that s dominates t , denoted $s \succeq t$, if $\sum_{j=1}^{\ell} s_j \geq \sum_{j=1}^{\ell} t_j$ for every $1 \leq \ell \leq n$. For any $0 \leq k \leq n$ we define the set $\mathcal{X}_{n,k} := \{x \in \{0, 1\}^n : \sum x_i = k\}$. Clearly $\succeq$ defines a partial order on $\mathcal{X}_{n,k}$, and we show now that it has a *small depth*.

Lemma 2 ([2]) *Let n, k be two integers with $0 \leq k \leq n$. Suppose we have a strictly increasing sequence*

$$b^0 \prec b^1 \prec \dots \prec b^q,$$

of vectors from $\mathcal{X}_{n,k}$. Then $q \leq k(n - k)$.

Proof: For each vector $\alpha \in \mathcal{X}_{n,k}$ we associate the number $\varphi(\alpha)$ defined by $\varphi(\alpha) = \sum_{\ell=1}^n \sum_{i=1}^{\ell} \alpha_i$.

If $\alpha \prec \beta$ then $\sum_{j=1}^{\ell} \alpha_j \leq \sum_{j=1}^{\ell} \beta_j$ for every $1 \leq \ell \leq n$ and the inequality is strict for at least one ℓ . We conclude that $\alpha \prec \beta$ implies $\varphi(\alpha) < \varphi(\beta)$.

Therefore the vectors with extreme values for φ are $\alpha = (0, \dots, 0, \underbrace{1, \dots, 1}_k)$ and $\beta = (\underbrace{1, \dots, 1}_k, 0, \dots, 0)$.

Since $\varphi(\alpha) = k(k - 1)/2$ and $\varphi(\beta) = k(k - 1)/2 + k(n - k)$, this concludes the proof. $\square$

A well-known characterization of the feasible instances of the 2-COLOR TOMOGRAPHY PROBLEM can be expressed using dominance.

Lemma 3 ([8]) *Let (r, s) be an instance of the 2-color tomography problem, such that r is non-increasing. Then (r, s) is feasible if and only if $r \preceq s^*$. Moreover if $r = s^*$, then there is a single solution, namely the realization having the first s_j cells of column j colored black, and the others white.*

There is a very simple graphical interpretation of this. Again let M be a matrix where in column j the first s_j cells are colored black and the remaining cells white. Then the row projection of M is s^* , and if $s^* = r$ we are done. Now if $s^* \neq r$, then some of the black cells in M have to be exchanged with some white cells in the same column but a lower row. These operations transform the matrix in such a way, that the new row projection is dominated by s^* . So if s^* does not dominate r , then there is no solution to the instance.

2 The gadget

The gadget depends on some integers n, k, u, v with $1 \leq k, u, v \leq n$ and $u \neq v$ as well as on two vectors $\alpha, \beta \in \mathcal{X}_{n,k}$. It is defined as the instance of n rows, and $2n+2$ columns with the following projections for $1 \leq i, j \leq n$

$$\begin{aligned} r_i^R &= \begin{cases} i+1 & \text{if } i \in \{u, v\} \\ i & \text{otherwise} \end{cases} & r_i^G &= \begin{cases} i & \text{if } i \in \{u, v\} \\ i+1 & \text{otherwise} \end{cases} \\ s_j^R &= n - j + \alpha_j & s_j^G &= 0 \\ s_{n+1}^R &= 1 & s_{n+1}^G &= n - 1 \\ s_{n+2}^R &= n - k + 1 & s_{n+2}^G &= k - 1 \\ s_{n+2+j}^R &= 0 & s_{n+2+j}^G &= n - j + 1 - \beta_j. \end{aligned}$$

Lemma 4 *If the instance above is feasible then $\alpha \preceq \beta$. Moreover, if $\alpha = \beta$ then the instance is feasible if and only if $\alpha_u + \alpha_v \geq 1$.*

Proof: Assume the instance is feasible, we will show that this implies $\alpha \preceq \beta$. Consider the yellow projection vectors $r^Y = 2n+2 - r^R - r^G$ and $s^Y = n - s^R - s^G$. We have that $r_i^Y = 2(n-i) + 1$ for $1 \leq i \leq n$. Note that r^Y is a non-increasing vector. Similarly, we obtain that $s_j^Y = j - \alpha_j$ and $s_{n+2+j}^Y = j - 1 + \beta_j$, for $1 \leq j \leq n$, and $s_{n+1}^Y = s_{n+2}^Y = 0$. The conjugate of the column yellow projections is a vector $(s^Y)^*$ with

$$(s^Y)_i^* = 2(n-i) + 1 - \alpha_i + \beta_i.$$

Then clearly $r^Y \preceq (s^Y)^*$ if and only if $\alpha \preceq \beta$. By assumption the 3-color instance $(r^R, r^G, r^Y, s^R, s^G, s^Y)$ is feasible, therefore the 2-color instance (r^Y, s^Y) is feasible as well — where yellow is renamed as black — which by Lemma 3 implies $r^Y \preceq (s^Y)^*$ and therefore also $\alpha \preceq \beta$. This shows the first part of the lemma.

Now assume that the instance has a solution, and $\alpha = \beta$. The $n \times (2n+2)$ grid is divided into 3 parts (see figure 2): into an $n \times n$ block (called *RY-block*), a $n \times 2$ rectangle (called *2-column translator*) and another $n \times n$ block (called *GY-block*). Again every block is sub-divided into an upper triangle, a diagonal and a lower triangle.

Since $\alpha = \beta$, we have $r^Y = (s^Y)^*$. So by Lemma 3 any solution must color in yellow the s_j^Y first cells in every column j , and no other cell. In particular it means that the lower triangle of the RY-block must be red, the lower triangle of the GY-block must be green, and both upper triangles have to be yellow.

Also on the first diagonal, the cell (i, i) has to be red if $\alpha_i = 1$ and yellow otherwise. On the second diagonal, the cell $(n+2+i, i)$ must be yellow if $\alpha_i = 1$ and green otherwise.




Figure 2: The structure of the gadget (left) and a realization (right) for $n = 7$, $k = 3$, $u = 2$, $v = 5$ and $\alpha = \beta = (0, 0, 1, 0, 1, 1, 0)$.

What can we say about the colors of the translator? If $\alpha_u = \alpha_v = 0$, then the cells $(n+1, u)$, $(n+2, u)$, $(n+1, v)$, $(n+2, v)$ have to be all red to satisfy the row projections. This contradicts the column projection $s_{n+1}^R = 1$, and hence the instance is not feasible.

Conversely, assume $\alpha_u + \alpha_v \geq 1$. We will color the cells of the translator in a manner that respects the required projections. If $i \notin \{u, v\}$ and $\alpha_i = 1$ — that is (i, i) is red — we color the cells $(n+1, i)$, $(n+2, i)$ in green. If $i \notin \{u, v\}$ and $\alpha_i = 0$, we color the cell $(n+1, i)$ in green and $(n+2, i)$ in red.

Without loss of generality assume that $\alpha_u = 1$. Hence (u, u) is red and we color $(n+1, u)$ in green and $(n+2, u)$ in red. We color $(n+1, v)$ in red. In addition we color $(n+2, v)$ in red if $\alpha_v = 0$ and in green otherwise. It can be verified that the coloring defined above is a solution to the instance, which concludes the proof of the lemma. $\square$

3 The reduction

In this section we will construct a reduction from VERTEX COVER to 3-COLOR TOMOGRAPHY. We basically use the same approach than in [2], but with a different gadget.

VERTEX COVER is a well known intractable problem, indeed one of the first 21 problems shown to be NP-complete by Karp [6].

VERTEX COVER PROBLEM

Input: a graph $G = (V, E)$ and an integer k .

Output: a set $S \subseteq V$ of size $|S| = k$ such that for every $(u, v) \in E$, $u \in S$ or $v \in S$.

Given an instance (G, k) of VERTEX COVER, we construct an instance (r^R, r^B, s^R, s^B) of the 3-COLOR TOMOGRAPHY PROBLEM which is feasible if and only if the former instance has a solution. Without loss of generality we assume that $k \leq n - 2$.

Let be $n = |V|$, $m = |E|$, and $N = k(n - k)(m - 1) + 1$. We denote the m edges as $E = \{e_0, e_1, \dots, e_{m-1}\}$, and the n vertices as $V = \{1, 2, \dots, n\}$. We define an instance with $N(n+1) + 1$ rows and $N(n+2) + n$ columns.

For row $p = 1, \dots, N(n+1) + 1$, let

$$x = \lfloor (p-1)/(n+1) \rfloor$$

$$i = (p-1) \bmod (n+1).$$

We think the set of rows as divided into N blocks of $n+1$ rows each, and a last block with a single row. We have x as the block index and i the row index relative to the block, with $0 \leq i \leq n$. Let

$t = x \bmod m$ and consider the edge $e_t = (u, v)$. We define the projections

$$r_p^R = x(n+2) + \begin{cases} n-k & \text{if } x < N \text{ and } i = 0 \\ 0 & \text{if } x = N \text{ and } i = 0 \\ i+1 & \text{if } i \in \{u, v\} \\ i & \text{if } i \in \{1, \dots, n\} \setminus \{u, v\} \end{cases}$$

$$r_p^G = (N-x-1)(n+2) + \begin{cases} n+2 & \text{if } x = 0 \text{ and } i = 0 \\ n+2+k & \text{if } x > 0 \text{ and } i = 0 \\ i & \text{if } i \in \{u, v\} \\ i+1 & \text{if } i \in \{1, \dots, n\} \setminus \{u, v\}. \end{cases}$$

In the same manner, for column $q = 1, \dots, N(n+2) + n$, let

$$y = \lfloor (q-1)/(n+2) \rfloor$$

$$j = ((q-1) \bmod (n+2)) + 1.$$

The reason for defining j this way, is that if cell (p, q) is part of an RY-block or an GY-block, then (i, j) will be the relative position inside the block with ranges $1 \leq i, j \leq n$. Similarly as for the rows, we think the set of columns as divided into N blocks with $n+2$ columns each and a last block with only n columns. Again, we have y as the block index, and j as the column index relative to a block with $1 \leq j \leq n+2$. For block $0 \leq y \leq N-1$ we define the red column projections as

$$s_q^R = (N-y-1)(n+1) + 1 + \begin{cases} n-j+1 & \text{if } j \in \{1, \dots, n\} \\ 1 & \text{if } j = n+1 \\ n-k+1 & \text{if } j = n+2, \end{cases}$$

For $y = N$ we set $s_q^R = 0$, for each $j = 1, \dots, n$. Similarly, for $y = 0$ the green column projections are

$$s_q^G = \begin{cases} 0 & \text{if } j \in \{1, \dots, n\} \\ n & \text{if } j = n+1 \\ k & \text{if } j = n+2. \end{cases}$$

and for $1 \leq y \leq N$ they are defined as

$$s_q^G = (y-1)(n+1) + 1 + \begin{cases} j & \text{if } j \in \{1, \dots, n\} \\ n-1 & \text{if } j = n+1 \\ k-1 & \text{if } j = n+2. \end{cases}$$

Clearly this is a polynomial time reduction. It remains to show the following theorem.

Theorem 1 *The 3-color tomography instance is feasible if and only if the vertex cover instance is feasible.*

Proof: For one direction of the statement, assume that the vertex cover instance is feasible, and let $b \in \mathcal{X}_{n,k}$ be the characteristic vector of a vertex cover of size k , i.e. $b_i = 1$ if and only if i belongs to the vertex cover.

We construct now a solution to the tomography instance. Consider the partitioning of the grid, as in figure 3. For convenience we refer to the source also as the 0-th row translator and to the sink as the $(N+1)$ -th row translator. The j -th cell of the x -th row translator is defined as $(x(n+1)+1, x(n+2)+j)$. We color the R-frame in red and the G-frame in green.

Let be any $1 \leq j \leq n$. We color the j -th cell of the source in yellow if $b_j = 1$ and in red otherwise. For $x = 1, \dots, N-1$ we color the j -cell of the x -th row translator in green if $b_j = 1$ and in red otherwise. In the sink we color the j -th cell in green if $b_j = 1$ and in yellow otherwise.

Now for block $x = 0, \dots, N-1$, consider the instance to the gadget defined by $\alpha = \beta = b$, and u, v such that $(u, v) = e_{x \bmod m}$. By Lemma 4 it is feasible, since b is a vertex cover and hence




Figure 3: The general structure of our reduction

$b_u + b_v \geq 1$. Then we color the $(n+1) \times (2n+2)$ cells starting at $((x-1)(n+2)+1, (x-1)(n+1)+1)$ exactly as in the solution to the gadget. It is straightforward to check that this grid satisfies the required projections, and therefore the tomography instance is feasible.

For the converse, assume that the tomography instance has a solution. For every $x = 1, \dots, N$ we apply Lemma 1 for the red color and intervals $I = [x(n+1)+1, N(n+1)+1]$ and $J = [1, x(n+2)]$. We deduce that in the solution the R-frame must be all red, and all GY-blocks (and also the G-frame) must be free of any red. Similarly, we show that the G-frame must be all green, and all RY-blocks must be free of any green.

This implies that in the source, k cells are yellow, and $n - k$ are red, in the row translators k cells are green and $n - k$ red, and in the sink k cells are green and $n - k$ yellow. We define the vectors $b^0, b^1, \dots, b^N \in \mathcal{X}_{n,k}$, such that for all $1 \leq j \leq n$ we have

- $b_j^0 = 1$ iff the j -th cell in the source is yellow,
- $b_j^x = 1$ iff the j -th cell in the x -th row translator is green, for all $1 \leq x \leq N$.

For $x = 0, \dots, N$, consider the part P of the solution that is the intersection of rows $[x(n+1)+2, x(n+1)+n+1]$ and columns $[x(n+2)+1, x(n+2)+2n+2]$. We number the rows of P from 1 to n and the columns from 1 to $2n+2$. Let $(u, v) = e_{x \bmod m}$. By subtracting from the row projections the number of red and green cells in the frames, we deduce that row $1 \leq i \leq n$ in P contains $i+1$ red cells and i green cells if $i \in \{u, v\}$ and i red cells and $i+1$ green cells if $i \notin \{u, v\}$.

We proceed similarly for the columns $n+1$ and $n+2$. By subtracting from the column projections the quantities that are in the frames, we deduce that column $n+1$ of P contains one red cell, and $n-1$ green cells, and column $n+2$ contains $n-k+1$ red cells and $k-1$ green cells.

Column $x(n+2)+j$ for $1 \leq j \leq n$ contains $n-j+1$ red cells that are not in the R-frame. Since GY-blocks are free of red, these cells must either be in the x -th row translator or in column j of P . Note that the j -cell of the x -th row translator is red iff $b_j^x = 0$. Therefore column j of P contains $n-j+b_j^x$ red cells and no green cell. Similarly column $n+2+j$ of P contains $n-j+1-b_j^{x+1}$ green cells and no red cell.

This implies that P is the solution to the gadget defined by u, v, α, β with $\alpha = b^x$ and $\beta = b^{x+1}$. Then by Lemma 4 we obtain that $b^x \preceq b^{x+1}$ and in general

$$b^0 \preceq b^1 \preceq \dots \preceq b^N.$$

By the choice of N and Lemma 2 there exists an ℓ such that

$$b^\ell = b^{\ell+1} = \dots = b^{\ell+m}.$$

By Lemma 4, we have $b_u^\ell + b_v^\ell \geq 1$ for all $(u, v) \in \{e_\ell, e_{\ell+1 \bmod m}, \dots, e_{\ell+m-1 \bmod m}\} = E$. Therefore b^ℓ encodes a vertex cover of size k , and this completes the proof. $\square$

4 Related problems

4.1 Edge-colored graphs with prescribed degrees

We can reduce the 3-COLOR TOMOGRAPHY PROBLEM to a similar graph problem.

Finding edge-colored graphs with prescribed degrees Let be a set of two colors $\{R, G\}$, and a vertex set V . We are given prescribed degrees $d^R, d^G : V \rightarrow \mathbb{N}$ and have to find two disjoint edge sets $E^R, E^G \subseteq V^2$ such that the graph $G(V, E^R \cup E^G)$ has the required degrees, i.e. for all $v \in V$

$$d^R(v) = |\{u : (u, v) \in E^R\}| \quad d^G(v) = |\{u : (u, v) \in E^G\}|.$$

Note that in contrast, finding an uncolored graph with given degree sequences can be solved in polynomial time, see for example [7].

Lemma 5 *The problem of finding an edge-colored graph with prescribed degrees is NP-hard.*

Proof: We reduce from the 3-color tomography problem. Let (r^R, r^G, s^R, s^G) be an $m \times n$ -instance of the 3-color tomography problem. We set $k = n + m$, $V = \{1, \dots, k\}$, and the following degrees, for $1 \leq i \leq n$ and $1 \leq j \leq m$

$$\begin{aligned} d^R(i) &= r_i^R + n - 1 & d^G(i) &= r_i^G \\ d^R(n+j) &= s_j^R & d^G(n+j) &= s_j^G + m - 1. \end{aligned}$$

Now we show that the instance (r^R, r^G, s^R, s^G) is feasible if and only if the instance (d^R, d^G) is feasible. For one direction, assume that there is a solution M to the 3-color tomography instance. We construct a solution E^R, E^G to the graph problem as follows. For any $1 \leq i \leq n$ and $1 \leq j \leq m$, if $M_{ij} = R$, then $(i, n+j) \in E^R$, if $M_{ij} = G$, then $(i, n+j) \in E^G$. Also for any $1 \leq i < i' \leq n$, we have $(i, i') \in E^R$ and for any $1 \leq j < j' \leq m$, we have $(n+j, n+j') \in E^G$. Now clearly E^R, E^G satisfy the required degrees.

For the converse, we define the quantity $\Phi = \sum_{i=1}^n d^R(i) - \sum_{j=1}^m d^R(n+j)$. By assumption (1) this value is $n(n-1)$. Since this value equals also

$$|E^R \cap \{1, \dots, n\}^2| - |E^R \cap \{n+1, \dots, n+m\}|,$$

there is a red edge between every pair of vertices (i, i') with $1 \leq i < i' < n$, and no edge between every pair of vertices $(n+j, n+j')$ with $1 \leq j < j' \leq m$. Similarly we can show that there is a green edge between every pair of vertices $(n+j, n+j')$ with $1 \leq j < j' \leq m$.

Now let M be the $m \times n$ grid, with cell (i, j) colored in red if $(i, n+j) \in E^R$, and in green if $(i, n+j) \in E^G$. By the degree requirements, M is a solution to the 3-color tomography instance. $\square$

4.2 Tiling Tomography

Tiling tomography was introduced in [3], and it consists of constructing a tiling that satisfies some given row and column projections for each type of tiles we admit.

Formally a tile is a finite set T of cells of the grid $\mathbb{N} \times \mathbb{N}$, that are 4-connected, in the sense that the graph $G(T, E)$ is connected for $E = \{((i, j), (i', j')) : |i - i'| + |j - j'| = 1\}$. By $T + (i', j') = \{(i + i', j + j') : (i, j) \in T\}$ we denote a copy of T that is shifted i' units down and j' units to the right. We say that a set of tiles is *feasible* if they do not intersect. In addition we say that it tiles the $m \times n$ grid if its (disjoint) union equals the set of all grid cells, and we refer it as a *tiling*.

In the tiling tomography problem we are given a finite set of tiles $\mathcal{T} = \{T_1, \dots, T_k\}$, and vectors $r^d \in \mathbb{N}^m, s^d \in \mathbb{N}^n$ for $1 \leq d \leq k$. The goal is to compute a matrix $M \in \{0, 1, \dots, k\}$ such that the set

$$\bigcup_{1 \leq d \leq k} \{T_d + (i, j) : 1 \leq i \leq n, 1 \leq j \leq m \text{ such that } M_{ij} = d, \text{ for } 1 \leq d \leq k\}$$

is a tiling of the $m \times n$ grid, with the projections

$$r_i^d = |\{j : M_{ij} = d\}| \quad s_j^d = |\{i : M_{ij} = d\}|.$$

By *width* and *height* of a tile T we understand the size of the smallest intervals I, J such that $T \subseteq I \times J$. This definition extends to set of tiles. A tile T is said to be rectangular if for every (i', j') such that $\{T, T + (i', j')\}$ is feasible, we have that the width of $\{T, T + (i', j')\}$ is at least twice the width of T or the height of the set is at least twice the height of T .

It was conjectured in [3], that for T_1 being a single cell and T_2 a non-rectangular tile, the $\{T_1, T_2\}$ -tiling tomography problem is NP-hard. This question is still open and intriguing.

4.3 Rectangular tiles

Consider two rectangular tiles, T_1 being a $p_1 \times q_1$ rectangle and T_2 a $p_2 \times q_2$ rectangle, i.e. $T_c = \{0, \dots, p_c - 1\} \times \{0, \dots, q_c - 1\}$, for $c \in \{1, 2\}$. What can be said about the complexity of the $\{T_1, T_2\}$ -tiling tomography problem?

If $\gcd(p_1, p_2) = d > 1$, then clearly any solution $\bar{M} \in \{0, 1, 2\}^{m \times n}$ to a $\{T_1, T_2\}$ -tiling tomography instance (r^c, s^c) , must satisfy that if $\bar{M}_{ij} \neq 0$, then $i \bmod d = 1$. Therefore the $\{T_1, T_2\}$ -tiling tomography problem can be reduced to the $\{T'_1, T'_2\}$ -tiling tomography problem, with T'_1 being a $(p_1/d) \times q_1$ rectangle, and T'_2 a $(p_2/d) \times q_2$ rectangle. We omit the formal reduction, which is straightforward.

From now on suppose that $\gcd(p_1, p_2) = \gcd(q_1, q_2) = 1$. We distinguish the following cases, up to row-column symmetry.

- If $p_1 = p_2 = 1$, that is the tiles are two horizontal bars of length q_1 and q_2 , then the problem can be solved in polynomial time (Theorem 2). We use an idea already present in [4], where it is proven for $q_1 = 1$.
- If $p_1 = q_2 = 1$ and $p_2 = q_1 = 2$, then the tiles are called dominoes, and again the problem can be solved in polynomial time, although with a more involved algorithm [9].
- If $p_1 = q_2 = 1, p_2 \geq 2$ and $q_1 \geq 3$ then the problem is open. The first author conjectures that the problem is NP-hard, while the other two conjecture that it could be solved in polynomial time with a similar approach as in [9].
- If $p_1, q_1 \geq 2$, then the problem is NP-hard (Theorem 3). In [3] the special case $p_1 = q_1 = 2, p_2 = q_2 = 1$ was related to the 3-color tomography problem, and it is therefore also NP-hard. We generalize this reduction in section 4.6
- If there is a third rectangular tile T_3 , then for the tile set $\{T_1, T_2, T_3\}$ the problem is NP-hard, see section 4.7.




Figure 4: Transforming a solution M for $c = 1$.

4.4 An algorithm for vertical bars

Theorem 2 *The tiling tomography problem can be solved in polynomial time for two rectangular tiles of dimensions $p_1 \times 1$ and $p_2 \times 1$.*

Proof: The algorithm is the simple greedy algorithm, as the one used in [4]. It iteratively *stacks* bars in the matrix.

Formally the algorithm is defined like this. We construct a matrix $A \in \{0, 1, 2\}^{m \times n}$ with the required projections. Initially A is all 0. We maintain a vector v such that v_j is the minimal i such that $A_{i,j} \neq 0$, and $v_j = m + 1$ if column j of A is all zero. Initially $v_j = m + 1$ for all $1 \leq j \leq n$. We also maintain vectors $\bar{r}^1, \bar{r}^2, \bar{s}^1, \bar{s}^2$, which represent the remaining projections. Initially they equal the given projections of the instance. The vectors $(v, \bar{r}^1, \bar{r}^2, \bar{s}^1, \bar{s}^2)$ define a more general tiling problem, where in every column j , only the first $v_j - 1$ cells have to be tiled.

The algorithm: Let $i = \max v_j$. If $i = 1$ we are done, and return A , if all vectors $\bar{r}^1, \bar{r}^2, \bar{s}^1, \bar{s}^2$ are zero, and return “no solution” otherwise.

If $i > 1$, let $i_1 = i - p_1$ and $i_2 = i - p_2$. If $\bar{r}_{i_1}^1 = \bar{r}_{i_2}^2 = 0$, abort and return “no solution”. Otherwise let $c \in \{1, 2\}$ such that $\bar{r}_{i_c}^c > 0$. Let j be a column with $v_j + 1 = i$ that maximizes $\bar{s}_j^c$. Then *drop* the bar $p_c \times 1$ in column j , i.e. set $A_{i_c,j} = c$, and decrease $\bar{r}_{i_c}^c$ and $\bar{s}_j^c$. Repeat the whole step.

Clearly, if this algorithm produces a matrix, then it defines a valid tiling with the required projections. We have to show that if the instance has a solution, then the algorithm will actually find one. For this purpose, let be some step of algorithm such that the intermediate instance $\mathcal{I} := (v, \bar{r}^1, \bar{r}^2, \bar{s}^1, \bar{s}^2)$ is feasible. The initial step could be a candidate. Let M be a solution to it. Let $i = \max v_j$. If $i = 1$, then $\bar{r}^1, \dots, \bar{s}^2$ are all zero, since the instance is feasible.

Let $i_1 = i - p_1$ and $i_2 = i - p_2$. We have that either $M_{i_1,j} = 1$ or $M_{i_2,j} = 2$ for every column j satisfying $v_j = i$, since M is a valid tiling. Therefore some of $\bar{r}_{i_1}^1, \bar{r}_{i_2}^2$ must be non zero. Let c, j be the values the algorithm chooses. Let $\mathcal{I}'$ be the instance obtained after the iteration of the algorithm, that is $\bar{r}_{i_c}^c, \bar{s}_j^c$ are decreased by 1 and v_j by p_c .

If $M_{i_c,j} = c$, then M' which equals M except for $M_{i_c,j} = 0$ is a solution to $\mathcal{I}'$.

If $M_{i_c,j} \neq c$, then by the projections, there must be another column k with $v_k = i$ and $M_{i_c,k} = c$. We will now transform M such that $M_{i_c,j} = c$. Then we are in the case above and done.

By the choice of the algorithm we have $\bar{s}_k^c \leq \bar{s}_j^c$. By this inequality, there exists i_0 such that the total number of c 's below the row i_0 is the same in both column j and column k . Take i_0 being the largest one satisfying that. By the choice of i_0 we have that $M_{i_0,j} = c$ and $M_{i_0,k} \neq c$. Since M is a valid tiling, then the restriction to cells below i_0 in column k is also a tiling and then $M_{i_0,k} \neq 0$. We conclude that between i_0 and i the number of 1's and 2's in column j is the same as in column k . Then exchanging the parts of columns j and k in M between i_0 and i , does not change the projections of M , and we obtain the required property $M_{i_c,j} = c$.

By the choice of the algorithm we have $s_k^c \leq s_j^c$. Now we claim that there is a row i_0 such between i_0 and i the number of 1's and 2's in column j is the same as in column k . Indeed, consider the largest i_0 such that the total number of c 's below the row i_0 is the same in both column j and column k . It must be that $M_{i_0,j} = c$ and $M_{i_0,k} \neq 0$ since M is a valid tiling. Then exchanging the parts of columns j and k in M between i_0 and i , does not change the projections of M , and we obtain the required property $M_{i_0,j} = c$. $\square$

4.5 A general NP-hardness proof structure

In the next section we will reduce the 3-color tomography problem to the tiling tomography problem for some fixed set of tiles $\mathcal{T}$. The proof uses a particular structure that we explain now.

Let $(r^R, r^G, r^Y, s^R, s^G, s^Y)$ be an instance to the 3-color tomography problem for an $m \times n$ grid. In the reduction we will choose constant size grid $\ell \times k$ — that we call a *block* — and three $\mathcal{T}$ -tilings of it, that we denote $\bar{M}^R, \bar{M}^G, \bar{M}^Y$. There will be two requirements: Let $\bar{r}^{c,d}, \bar{s}^{c,d}$ be the T_d -projections of the tiling $\bar{M}^c$ for $c \in \{R, G, Y\}$ and $d \in \{1, 2\}$.

The first requirement is that the vectors $\{\bar{r}^{R,1}, \bar{r}^{G,1}, \bar{r}^{Y,1}\}$ are affine linear independent. The same requirement holds for the column projections $\{\bar{s}^{R,1}, \bar{s}^{G,1}, \bar{s}^{Y,1}\}$. This implies that every vector r spanned by $\bar{r}^{R,1}, \bar{r}^{G,1}, \bar{r}^{Y,1}$, has a unique decomposition into $r = n_R \bar{r}^R + n_G \bar{r}^G + n_Y \bar{r}^Y$ for $n_R + n_G + n_Y = n$.

The reduction, consists of an $m\ell \times nk$ grid, and the projections $1 \leq i \leq \ell, 1 \leq j \leq k, 1 \leq x \leq m, 1 \leq y \leq n, d \in \{1, 2\}$

$$\begin{aligned} r_{x\ell-\ell+i}^d &= \sum_c r_x^c \cdot \bar{r}_i^{c,d} \\ s_{yk-k+j}^d &= \sum_c s_y^c \cdot \bar{s}_j^{c,d} \end{aligned}$$

The idea is that the $m\ell \times nk$ is partitioned into mn blocks of dimension $\ell \times k$. **The second requirement** is that in every solution $\bar{M}$ to the tiling instance, all blocks of $\bar{M}$, are either $\bar{M}^R, \bar{M}^G, \bar{M}^Y$ or blocks that have equivalent projections.

Lemma 6 *The instance to the $\mathcal{T}$ -tiling problem has a solution if and only if the instance to the 3-color tomography problem has a solution.*

Proof: Let $M \in \{R, G, Y\}^{m \times n}$ be a solution to the 3-color tomography problem. We transform it into a matrix $\bar{M} \in \{0, 1, 2\}^{m\ell \times nk}$ by replacing each cell (i, j) of M by the $\ell \times k$ matrix $\bar{M}^c$ for $c = M_{ij}$. By construction, this is a solution to the tiling problem.

For the converse, suppose that there is a solution $\bar{M}$ to the tiling problem. By the second requirement, every block of $\bar{M}$ can be associated to one of the colors $\{R, G, Y\}$. We construct a matrix $M \in \{R, G, Y\}^{m \times n}$ such that $M_{xy} = c$ if the block (x, y) of $\bar{M}$ is $\bar{M}^c$, or something projection equivalent.

Fix some arbitrary $1 \leq x \leq m$. By the first requirement, the projections of the rows $x\ell - \ell + 1, \dots, x\ell$ have a unique decomposition into $n_R r^{R,1} + n_G r^{G,1} + n_Y r^{Y,1}$ with $n_R + n_G + n_Y = n$. By the definitions of the projections $n_R = r_x^R, n_G = r_x^G, n_Y = r_x^Y$, and then row x of M has the required projections. We proceed in the same manner for the columns and show that M is a solution to the 3-color tomography instance. $\square$

4.6 An NP-hardness proof for two rectangular tiles

Theorem 3 *The tiling tomography problem is NP-hard for two rectangular tiles of dimensions $p_1 \times q_1$ and $p_2 \times q_2$ with $\gcd(p_1, p_2) = \gcd(q_1, q_2) = 1$ and $p_1, q_1 \geq 2$.*




Figure 5: The 3 valid block tilings.

Proof: We apply Lemma 6 for $\ell = 2p_1p_2$ and $k = 2q_1q_2$. The 3 tilings of the $\ell \times k$ grid are depicted in figure 5, and defined formally as follows. The rows $I = \{1, \dots, \ell\}$ and the columns $J = \{1, \dots, k\}$ are partitioned into sets I_1, I_2, I_3, I_4 and J_1, J_2, J_3, J_4 defined as

$$\begin{aligned}
 I_1 &= \{1, \dots, p_2\} & J_1 &= \{1, \dots, q_2\} \\
 I_2 &= \{p_2 + 1, \dots, p_1p_2\} & J_2 &= \{q_2 + 1, \dots, q_1q_2\} \\
 I_3 &= \{p_1p_2 + 1, \dots, p_1p_2 + p_2\} & J_3 &= \{q_1q_2 + 1, \dots, q_1q_2 + q_2\} \\
 I_4 &= \{p_1p_2 + p_2 + 1, \dots, 2p_1p_2\} & J_4 &= \{q_1q_2 + q_2 + 1, \dots, 2q_1q_2\}.
 \end{aligned}$$

Then $\bar{M}^R$ is defined as the block tiling that covers $(I_1 \cup I_4) \times (J_3 \cup J_4)$ with T_2 and the rest with T_1 , $\bar{M}^G$ is defined as the block tiling that covers $(I_3 \cup I_4) \times (J_1 \cup J_4)$ with T_2 and the rest with T_1 , while $\bar{M}^Y$ is defined as a tiling using only T_1 . These tilings are uniquely defined. Clearly the row T_1 -projections of the 3 tilings are affine linear independent, so the first requirement of the construction is satisfied.

The second requirement follows from a sequence of observations. Let $\bar{M}$ be the solution to the tiling instance, obtained by reduction from a 3-color instance $(r^R, r^G, r^Y, s^R, s^G, s^Y)$.

First note that in the tilings $\bar{M}^R, \bar{M}^G, \bar{M}^Y$, every tile is completely contained in the $\ell \times k$ block. Therefore the tiling instance has zero projections for T_1 at rows x with $(x-1) \bmod \ell > \ell - p_1 + 2$. A similar observation holds for tile T_2 and for the column projections. As a result in $\bar{M}$ every tile is completely contained in some $\ell \times k$ block, and in other words every block of $\bar{M}$ is $\{T_1, T_2\}$ -tiling.

What can we say about the possible tilings? Again note that in the tilings $\bar{M}^R, \bar{M}^G, \bar{M}^Y$, every row in I_2 is completely covered by T_1 -tiles. Therefore by the projections, this holds also for every block in $\bar{M}$. The same observation can be done about columns in J_2 .

Note that if $ap_1 + bp_2 = 2p_1p_2$, then $(a, b) \in \{(0, 2p_1), (2p_2, 0), (p_2, p_1)\}$. This is simply because by $\gcd(p_1, p_2) = 1$, in any solution to $ap_1 = p_2(2p_1 - b)$, a must be a multiple of p_2 . Together with the previous observation, this implies that every column of a block is either covered completely by T_1 -tiles or covered half by T_1 -tiles and half by T_2 -tiles. The same observation holds for the rows.

The trickiest observation of this proof is that in every block of $\bar{M}$, the region $I_1 \times J_1$ is covered by T_1 . For a proof by contradiction, suppose it is covered by T_2 , in fact by a single tile T_2 since $|I_1 \times J_1| = |T_2|$. But since $I_2 \times J$ is covered with T_2 , and by $\gcd(q_1, q_2) = 1$, it must be that the cell $(p_2 + 1, q_2 + 1)$ is covered by a tile $T_2 + (p_2 + 1, j)$ for some column $j \leq q_2$. By the same argument, the cell $(p_2 + 1, q_2 + 1)$ is *also* covered by a tile $T_2 + (q_2 + 1, i)$ for some row $i \leq p_2$. Therefore these two tiles overlap in $(p_2 + 1, q_2 + 1)$, which contradicts that $\bar{M}$ is a (valid) tiling.

Now fix a block of $\bar{M}$. If row 1 is partly covered by T_2 , then T_2 -tiles must cover the half columns in J . Hence in the row 1 they cover exactly the columns in $J_3 \cup J_4$. The same argument shows that every column $j \in J_3 \cup J_4$ is then half covered by T_2 -tiles. Previous observation state that $I_2 \times \{j\}$ is covered by T_1 . But the length of I_2 is not a multiple of p_1 . Therefore $(p_1p_2 + 1, j)$ must then also be covered by T_1 and hence $(I_2 \cup I_3) \cup \{j\}$ is covered by T_1 -tiles. Therefore $(I_1 \cup I_4) \times \{j\}$ is covered by T_2 . The choice of j was arbitrary, and therefore the block-tiling is exactly $\bar{M}^R$.

Similarly we deduce that if column 1 is covered partly by T_2 , then the block-tiling is exactly $\bar{M}^G$. Now if row 1 and column 1 are completely covered by T_1 , then $(I_1 \cup I_2) \times J$ and $I \times (J_1 \cup J_2)$ are completely covered by T_2 -tiles. As a result the block-tiling only contains in $(I_3 \cup I_4) \times (J_3 \cup J_4)$ either T_1 -tiles or T_2 -tiles, that correspond with the $\bar{M}^Y$ tiling and another we call the bad tiling, respectively.

We will show that no bad tiling appears in $\bar{M}$. Let N_R be the number of blocks in $\bar{M}$ that are $\bar{M}^R$. Similarly, let N_B be number of bad block-tilings in $\bar{M}$. Note that the row projection of a bad tiling equal the row projections of $\bar{M}^G$ and that the column projections equal the projections of $\bar{M}^R$. Therefore by the projections we have the equalities

$$N_R = \sum_i r_i^R \quad N_R + N_B = \sum_j s_j^R.$$

Since by assumption $\sum_i r_i^R = \sum_j s_j^R$, we have $N_B = 0$. This shows the second requirement of our construction, and by Lemma 6 completes the proof. $\square$

4.7 An NP-hardness proof for three rectangular tiles

Theorem 4 *The tiling tomography problem is NP-hard for any 3 rectangular tiles.*

Proof:[sketch] Let $p_1 \times q_1$, $p_2 \times q_2$ and $p_3 \times q_3$ the respective dimensions of 3 tiles T_1, T_2, T_3 .

The idea of the construction is that we apply the general proof scheme from section 4.5 with 3 tilings $\bar{M}^R, \bar{M}^G, \bar{M}^Y$, such that $\bar{M}^R$ contains tile T_1 in position $(0, 0)$, $\bar{M}^G$ contains T_2 and $\bar{M}^Y$ contains T_3 in position $(0, 0)$. Moreover each of the 3 tiling minimizes lexicographically n_1, n_2, n_3 , where n_c is the number of tiles T_c in the tiling.

Formally, let i_1 be the smallest number $i > 0$ with $i \bmod p_1 = 0$ and either $i \bmod p_2 = 0$ or $i \bmod p_3 = 0$. Let i_2 be the smallest number $i \geq i_1$ with $(i - i_1) \bmod p_2 = 0$ and $i \bmod p_3 = 0$. Let i_3 be the smallest number $i > 0$ with $i \bmod p_2 = 0$ and $i \bmod p_3 = 0$. We define numbers j_1, j_2, j_3 in exactly the same manner with q_1, q_2, q_3 playing the same role as p_1, p_2, p_3 .

We apply Lemma 6 for $k = \max\{i_2, i_3\}$ and $\ell = \max\{j_2, j_3\}$. The 3 tilings of the $\ell \times k$ grid are depicted in figure 6, and defined formally as follows. In this section we assume for convenience that the rows and column indices relative to a block start at 0 instead of 1.

In $\bar{M}^R$, the $[1, i_1] \times [1, j_1]$ subsquare is completely tiled with T_1 . Then the region $[1, i_2] \times [1, j_2] - [1, i_1] \times [1, j_1]$ is completely tiled with T_2 and the remaining part with T_3 . Note that by the choice of k , no column intersects a tile T_1, T_2 and T_3 . For example column 0 intersects T_1 tiles from row 0 to $i_1 - 1$, and then either tiles T_3 from row i_1 to k (if $i_1 \bmod p_3 = 0$) or tiles T_3 from row i_1 to k (if $i_1 \bmod p_3 \neq 0$). The same holds for rows.

In $\bar{M}^G$, the $[1, i_3] \times [1, j_3]$ subsquare is completely tiled with T_2 , and the remaining part with T_3 . In $\bar{M}^Y$, the whole $[1, k] \times [1, \ell]$ block is tiled with T_3 .

Clearly the projections of these tiles satisfy the first requirement for the general proof structure. Now there are different interesting observations to make. In the tilings above, every tile $T_c + (i, j)$ satisfies $i \bmod p_c = 1$ and $j \bmod q_c = 1$. This means that the projections of tile T_c are zero for any row $i \neq 1 \pmod{p_c}$ or column $j \neq 1 \pmod{q_c}$, and in any solution M to the tiling instance resulting from the reduction, the property must hold for all blocks as well. This observation is crucial for the proof.

In particular it implies the following fact (*). Fix some solution M to the tiling instance resulting from the reduction. Consider a block in M . If there is some *vertical separation* between two types of tiles, in the sense that cell $(i - 1, j)$ is covered by some tile T_a and (i, j) by some tile T_b with $a \neq b$, then we must have $i \bmod p_a = 1$ and $i \bmod p_b = 1$.

We use this observation to show that the construction satisfies also the second requirement. Fix some solution M to the tiling instance resulting from the reduction. We distinguish 3 types of blocks: (1) Blocks that contain a tile T_1 , (2) blocks that do not contain any tile T_1 , but contain a tile T_2 , and (3) blocks that are completely tiled with T_3 . So blocks of the third type are exactly $\bar{M}^Y$, and we have to show that blocks of the first type are exactly $\bar{M}^R$ and blocks of the second type are $\bar{M}^G$.




Figure 6: The 3 valid block tilings.

Let n_R be the number of blocks of the first type, and consider one of them. Then by the projections all tiles T_1 must be contained in the region $[1, i_1] \times [1, j_1]$, and by the observation (*) above, the whole region must be tiled with T_1 . Now if $i_2 > i_1$, then the region $[1, i_2] \times [1, j_2] - [1, i_1] \times [1, j_1]$ cannot contain tiles T_1 nor T_3 and must be completely tiled with T_2 . Later we will show that the remaining part of the block is tiled with T_3 .

Note that n_R is also the total number of red projections in the original 3-color tomography instance, so if $i_3 \leq i_2$, then all T_2 -tiles that have to be placed in a block-row $i \geq \max\{i_2, i_3\}$, are placed in a type 1 block. The same observation can be made for columns. Therefore all tiles T_2 in a type 2 block, must be placed at positions of the form $(i, j) \in [1, i_3] \times [1, j_3]$. By the observation (*), the whole region $[1, i_3] \times [1, j_3]$ is completely tiled with T_2 . By the observation above, the remaining part can only be tiled with T_3 , which shows that the type 2 blocks are exactly $\bar{M}^G$.

Let n_G be the number of type 2 blocks in M . It is also the total number of green projections in the original 3-color tomography instance. Therefore by the projections, every tile $T_2 + (i, j)$ with $i > \max\{i_2, i_3\}$ or $j > \max\{j_2, j_3\}$ must be contained in a type 2 block. This shows that the remaining part $[1, k] \times [1, \ell] - [1, i_2] \times [1, j_2]$ of a type 1 block, contains only T_3 -tiles. This shows that the type 1 blocks are exactly $\bar{M}^R$.

Therefore the construction satisfies the second requirement for Lemma 6, and we are done. $\square$

5 Acknowledgement

We thank Christophe Picouleau and Dominique de Werra for correcting an earlier version of this manuscript.

References

- [1] A. Alpers, L. Rodek, H.F. Poulsen, E. Knudsen, and G.T. Herman. *Advances in Discrete Tomography and Its Applications*, chapter Discrete Tomography for Generating Grain Maps of Polycrystals, pages 271–301. Birkhäuser Boston, 2007.
- [2] M. Chrobak and C. Dürr. Reconstructing polyatomic structures from discrete X-rays: NP-completeness proof for three atoms. *Theoretical Computer Science*, 259:81–98, 2001.
- [3] Marek Chrobak, Peter Couperus, Christoph Dürr, and Gerhard Woeginger. On tiling under tomographic constraints. *Theoret. Comput. Sci.*, 290(3):2125–2136, 2003.

- [4] Christoph Dürr, Eric Goles, Ivan Rapaport, and Eric Rémila. Tiling with bars under tomographic constraints. *Theoret. Comput. Sci.*, 290(3):1317–1329, 2003.
- [5] R. Gardner, P. Gritzmann, and D. Prangenberg. On the computational complexity of reconstructing lattice sets from their X-rays. *Discr. Math.*, 202:45–71, 1999.
- [6] M.R. Garey and D.S. Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. W.H.Freeman and Co., 1979.
- [7] Attila Kuba and Gabor T. Herman. *Discrete tomography: Foundations, Algorithms and Applications*, chapter Discrete tomography: A Historical Overview. Birkhäuser, 1999.
- [8] H.J. Ryser. Matrices of zeros and ones. *Bull. Am. Math. Soc.*, 66:442–464, 1960.
- [9] Nicolas Thiant. *Constructions et reconstructions de pavages de dominos*. PhD thesis, Université Paris 6, 2006.